

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
17 May 2001 (17.05.2001)

PCT

(10) International Publication Number  
**WO 01/34563 A1**

(51) International Patent Classification<sup>7</sup>: C07C 315/06, 317/28

(21) International Application Number: PCT/US00/41609

(22) International Filing Date: 25 October 2000 (25.10.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/161,884 27 October 1999 (27.10.1999) US

(71) Applicant (for all designated States except US): NOBEX CORPORATION [US/US]; P.O. Box 13940, Research Triangle Park, NC 27709-3940 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): EKWURIBE, Nnochiri, N. [NG/US]; 216 Coltsgate Drive, Cary, NC 27511 (US). JAMES, Kenneth, D. [US/US]; 1607 Quaker Creek Drive, Mebane, NC 27302 (US).

(74) Agent: STRICKLAND, J., Michael; Myers Bigel Sibley & Sajovec, P.O. Box 37428, Raleigh, NC 27627 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: RESOLUTION OF INTERMEDIATES IN THE SYNTHESIS OF SUBSTANTIALLY PURE BICALUTAMIDE



of up to 6 carbons; or R<sup>4</sup> is phenyl which bears one, two or three substituents independently selected from hydrogen, halogen, nitro, carboxy, carbamoyl and cyano, and alkyl, alkoxy, alkanoyl, alkylthio, alkylsulphinyl, alkylsulphonyl, perfluoroalkyl, perfluoroalkylthio, perfluoroalkylsulphinyl, perfluoroalkylsulphonyl, alkoxycarbonyl and N-alkylcarbamoyl each of up to 4 carbon atoms, and phenyl, phenylthio, phenylsulphinyl and phenylsulphonyl; or R<sup>4</sup> is naphthyl; or R<sup>4</sup> is a 5- or 6-membered saturated or unsaturated heterocyclic which contains one, two or three heteroatoms selected from oxygen, nitrogen and sulfur, which heterocyclic may be a single ring or may be fused to a benzo-ring, and which heterocyclic is unsubstituted or bears one or two halogen, cyano or amino, or alkyl, alkoxy, alkylthio, alkylsulphinyl or alkylsulphonyl each of up to 4 carbon atoms, or oxy or hydroxy substituents, or which if sufficiently saturated may bear one or two oxo substituents; and X<sup>1</sup> is oxygen, sulfur, sulphinyl (-SO-), sulphonyl (-SO<sub>2</sub>-), imino (-NH-) or alkylimino (-NR<sup>5</sup>-) where R<sup>5</sup> is alkyl having up to 6 carbon atoms. The resolved intermediate compound of Formula (I) is then treated under conditions sufficient to provide a substantially pure enantiomer of an acylanilide such as Casodex® (bicalutamide) and/or its derivatives.

(57) Abstract: A method of preparing a substantially pure enantiomer of an acylanilide such as Casodex® (bicalutamide) and/or its derivatives includes resolving an intermediate compound having the structure of Formula (I), wherein: R<sup>1</sup> is alkyl or haloalkyl having up to 4 carbons; R<sup>2</sup> is alkyl having up to 6 carbon atoms; R<sup>3</sup> is a direct link or alkyl having up to 6 carbon atoms; R<sup>4</sup> is alkyl, alkenyl, hydroxyalkyl or cycloalkyl each

## RESOLUTION OF INTERMEDIATES IN THE SYNTHESIS OF SUBSTANTIALLY PURE BICALUTAMIDE

### Related Applications

This application claims priority from N. Ekwuribe, United States Provisional Application 60/161,884, filed October 27, 1999, the disclosure of which is incorporated by reference herein in its entirety.

5

### Field Of The Invention

The present invention relates to methods of resolving optically active organic compounds, and more particularly to methods of resolving optically active pharmaceutical compounds.


10

### Background Of The Invention

Androgen deprivation is a common treatment for persons with prostate cancer. Various non-steroidal antiandrogens are known for use in the treatment of prostate cancer. For example, bicalutamide, which may be among the most commonly used non-steroidal antiandrogens in the world, is typically used in the treatment of prostate cancer.

15 Bicalutamide is commercially available as Casodex<sup>®</sup> (bicalutamide) from Astra Zeneca Pharmaceuticals.

The chemical name of bicalutamide is N-[4-cyano-3-(trifluoromethyl)phenyl]-3-[(4-fluorophenyl)sulfonyl]-2-hydroxy-2-methyl-propanamide,(+-). The structural formula of bicalutamide is:



The  $\beta$ -carbon atom in the propanamide is a chiral carbon. As a result, bicalutamide is an optically active compound.


Optically active compounds have the ability to rotate the plane of polarized light. In describing an optically active compound, the prefixes D and L or R and S are used to denote the absolute configuration of the molecule about its chiral center(s). The prefixes d and l or (+) and (-) are used to denote the optical rotation of the compound (*i.e.*, the direction in which a plane of polarized light is rotated by the optically active compound). The l or (-) prefix indicates that the compound is levorotatory (*i.e.*, rotates the plane of polarized light to the left or counterclockwise) while the d or (+) prefix means that the compound is dextrarotatory (*i.e.*, rotates the plane of polarized light to the right or clockwise). The sign of optical rotation, (-) and (+), is not related to the absolute configuration of the molecule, R and S.

Optically active compounds, such as bicalutamide, exist as a pair of stereoisomers that are identical with the notable exception that they are non-superimposable mirror images of one another. A specific stereoisomer, such as the R isomer, may be referred to as an enantiomer. A mixture of R and S enantiomers may be referred to as a racemic mixture.

Bicalutamide, is presently commercially available as a racemic mixture. The racemic mixture of bicalutamide may be synthesized by various methods including, for example, the methods described in U.S. Patent No. 4,636,505 to Tucker. Tucker further describes various derivatives and analogs of bicalutamide having antiandrogenic properties. Additionally, Tucker states that this racemic mixture of acylanilide derivatives may be resolved into optically active forms which possess antiandrogenic activity. This method requires complete synthesis of the drug in a racemic mixture followed by esterification, resolution of the diastereomers and hydrolysis to obtain the desired enantiomer. Tucker fails to propose specific methods for achieving this resolution.


U.S. Patent No. 5,985,868 to Gray proposes synthesizing racemic mixtures of Casodex<sup>®</sup> (bicalutamide) using methods as described in U.S. Patent No. 4,636,505 to Tucker, and obtaining the (R)-(-) enantiomer of Casodex<sup>®</sup> (bicalutamide) by resolution of the enantiomers of Casodex<sup>®</sup> (bicalutamide) using fractional crystallization or chromatography

of diastereomeric esters of chiral acids. Gray notes that other standard methods of resolution such as simple crystallization and chromatographic resolution can also be used. The methods of Gray require complete synthesis of the drug in a racemic mixture. The racemic mixture of the drug is then modified to facilitate resolution, and modified again to yield the active enantiomeric compound. Gray further states that intermediates of Casodex<sup>®</sup> (bicalutamide) may be resolved using fractional crystallization or chromatography of diastereomeric esters of chiral acids. However, Gray fails to propose specific intermediates, specific methods, specific esters, and/or specific chiral acids for accomplishing such a resolution. Gray further states that a carboxylic acid precursor, 3-(4-fluorophenyl)-2-hydroxy-2-methylpropanoic acid, which has the following structure:




may be resolved by fractional crystallization of diastereomeric salts with chiral amines. However, Gray fails to propose specific methods, specific diastereomeric salts, and/or specific chiral amines for accomplishing such a resolution.


In Howard Tucker et al., *Resolution of the Nonsteroidal Antiandrogen 4'-Cyano-3-[(4-fluorophenyl)sulfonyl]-2-hydroxy-2-methyl-3'-(trifluoromethyl)-propionanilide and the Determination of the Absolute Configuration of the Active Enantiomer*, 31 J. MED. CHEM. 885-887 (1988), the authors propose preparing chiral bicalutamide by resolution of the thioether:



Once resolved, the enantiomers of the thioether may be oxidized to the sulfone by known means. The authors also propose resolution of the thioether by reaction of the thioether with (R)-(-)-camphanoyl chloride in pyridine to provide the diastereomeric ester:



The diastereomeric esters may then be separated by careful flash chromatography on silica gel. The individual pure diastereomeric isomers may then be hydrolyzed, without racemization, using methanolic sodium hydroxide to yield enantiomeric alcohols:




wherein  $E^1$  is  $\text{CH}_3$  and  $E^2$  is  $\text{OH}$  for the S configuration, or  $E^1$  is  $\text{OH}$  and  $E^2$  is  $\text{CH}_3$  for the R configuration. These methods require complete synthesis of the drug in a racemic mixture. The racemic mixture of the drug is then modified to facilitate resolution, and modified again to yield the active enantiomeric compound.

Synthesis of the entire drug prior to resolution may result in time and labor costs required for the performance of extra steps, and in the inefficient use of costly starting materials that become components of the less preferred S-enantiomer, which may be disposed of or recycled, resulting in even more expense. Consequently, there is a need in the art for a more streamlined method for preparing substantially enantiomerically pure bicalutamide, which eliminates these additional steps, and which makes efficient use of the starting materials and minimizes waste.

### Summary Of The Invention


Embodiments of the present invention provide a synthetic method comprising a resolution step which takes place prior to the addition of the most expensive components of the active compound. By resolving an intermediate compound rather than resolving the completely synthesized drug as described above for conventional methods, methods

according to the present invention may reduce or eliminate the need for additional post-synthesis procedures and/or reduce or eliminate the need to recycle the less-preferred enantiomer. Additionally, the use of expensive starting materials such as:



- 5 to produce the less-preferred enantiomer may be avoided, which may reduce the costs of producing a substantially pure form of the more-preferred enantiomer.

According to embodiments of the present invention, methods of preparing a substantially pure enantiomer of an acylanilide such as Casodex® (bicalutamide) and/or its derivatives are provided. The methods include resolving an intermediate compound having  
10 the structure of Formula I:



wherein


- $\text{R}^1$  is alkyl or haloalkyl having up to 4 carbons;  
 $\text{R}^2$  is alkyl having up to 6 carbon atoms;  
 15  $\text{R}^3$  is a direct link or alkyl having up to 6 carbon atoms;  
 $\text{R}^4$  is alkyl, alkenyl, hydroxyalkyl or cycloalkyl each of up to 6 carbons; or  $\text{R}^4$  is phenyl which bears one, two or three substituents independently selected from hydrogen, halogen, nitro, carboxy, carbamoyl and cyano, and alkyl, alkoxy, alkanoyl, alkylthio, alkylsulphinyl, alkylsulphonyl, perfluoroalkyl, perfluoroalkylthio,  
 20 perfluoroalkylsulphinyl, perfluoroalkylsulphonyl, alkoxycarbonyl and N-alkylcarbamoyl each of up to 4 carbon atoms, and phenyl, phenylthio, phenylsulphinyl and phenylsulphonyl; or  $\text{R}^4$  is naphthyl; or  $\text{R}^4$  is a 5- or 6-membered saturated or unsaturated heterocyclic which contains one, two or three heteroatoms selected from oxygen, nitrogen and sulfur, which heterocyclic may be a single ring or  
 25 may be fused to a benzo-ring, and which heterocyclic is unsubstituted or bears one or two halogen, cyano or amino, or alkyl, alkoxy, alkylthio, alkylsulphinyl or alkylsulphonyl each of up to 4 carbon atoms, or oxy or hydroxy substituents, or which if sufficiently saturated may bear one or two oxo substituents; and

$X^1$  is oxygen, sulfur, sulphinyl (-SO-), sulphonyl (-SO<sub>2</sub>-), imino (-NH-) or alkylimino (-NR<sup>5</sup>-) where R<sup>5</sup> is alkyl having up to 6 carbon atoms.

The resolved intermediate compound of Formula I is then treated under conditions sufficient to provide a substantially pure enantiomer of the acylanilide.

- 5 In some embodiments of the present invention, the step of resolving an intermediate compound of Formula I includes crystallizationally resolving the intermediate compound of Formula I. The crystallizationally resolving step includes contacting the intermediate compound of Formula I with a chiral base to provide a diastereomeric mixture of a chiral salt, crystallizationally resolving the diastereomeric mixture of the chiral salt to provide a
- 10 substantially pure enantiomer of the chiral salt, and recovering a substantially pure enantiomer of the intermediate compound of Formula I. The contacting step preferably includes contacting the intermediate compound of Formula I with (-)-cinchonidine to provide a diastereomeric mixture of the chiral salt. The step of crystallizationally resolving the diastereomeric mixture of the chiral salt preferably includes contacting the diastereomeric
- 15 mixture of the chiral salt with a solvent system comprising, for example, methylene chloride and diethyl ether. In other embodiments, the step of resolving an intermediate compound of Formula I may include high performance liquid chromatographically resolving the intermediate compound of Formula I.

- The resolved intermediate is preferably contacted with an aniline compound having
- 20 the structure of Formula II:



wherein

- R<sup>6</sup> is cyano, carbamoyl, nitro, fluoro, chloro, bromo, iodo, or hydrogen, or alkyl, alkoxy, alkanoyl, alkylthio, alkylsulphinyl, alkylsulphonyl, perfluoroalkyl, perfluoroalkylthio, perfluoroalkylsulphinyl or perfluoroalkylsulphonyl each having up to 4 carbon atoms,
- 25 or phenylthio, phenylsulphinyl or phenylsulphonyl;
- R<sup>7</sup> is cyano, carbamoyl, nitro, fluoro, chloro, bromo or iodo, or alkanoyl, alkylthio, alkylsulphinyl, alkylsulphonyl, perfluoroalkyl, perfluoroalkylthio,

perfluoroalkylsulphinyl or perfluoroalkylsulphonyl each of having up to 4 carbon atoms; or phenylthio, phenylsulphinyl or phenylsulphonyl; and

R<sup>8</sup> is hydrogen or halogen;

under conditions sufficient to provide a substantially pure enantiomer of an acylanilide. The  
5 substantially pure enantiomer of the acylanilide is preferably the (R)-enantiomer of Casodex<sup>®</sup> (bicalutamide).

By resolving intermediates rather than resolving completely synthesized drugs, methods according to the present invention may provide more cost effective ways of synthesizing substantially pure enantiomers of acylanilides such as Casodex<sup>®</sup> (bicalutamide)  
10 and derivatives thereof than are provided by conventional resolution techniques. For example, methods according to the present invention may be more cost effective because they resolve the intermediates prior to reacting them with the expensive aniline component, avoiding unwanted costs associated with the expensive aniline component, which is typically wasted in the production of the less-preferred enantiomer.

15

#### **Brief Description of the Drawings**

Figure 1 shows a <sup>19</sup>F NMR spectrum of the pure salt of the (R)-hydroxyacid and (-)-cinchonidine.

Figure 2 shows a <sup>19</sup>F NMR spectrum of the salt of the racemic (R,S)-hydroxyacid and  
20 (-)-cinchonidine prior to resolution.

Figure 3 shows a <sup>19</sup>F NMR spectrum of the salt of the racemic (R,S)-hydroxyacid and (-)-cinchonidine after one round of crystallization.

#### **Detailed Description Of Preferred Embodiments**


25 The invention will now be described with respect to preferred embodiments described herein. It should be appreciated however that these embodiments are for the purpose of illustrating the invention, and are not to be construed as limiting the scope of the invention as defined by the claims. As the substituents R<sup>1</sup>-R<sup>8</sup> and X<sup>1</sup> have been defined above, they will not be further defined herein other than to describe preferred substituents for the preferred  
30 embodiments. Unless otherwise noted, all percentages used herein are percent by weight. As used herein, the term "substantially enantiomerically pure" refers to a substance that has preferably between about 95% and 100% of one form (either R or S) and between about 5%



and 0% of the other form, more preferably between about 99% and 100% of one form (either R or S) and between about 1% and 0% of the other form, and, most preferably, between about 99.9% and 100% of one form (either R or S) and about 0.1% and 0% of the other form.

Embodiments of the present invention provide methods of preparing a substantially  
 5 pure enantiomer of an acylanilide. Particularly preferred methods according to the present invention resolve a racemic mixture of intermediate compounds to provide Casodex<sup>®</sup> (bicalutamide) and/or its derivatives in a more cost effective manner than conventional methods that resolve the racemic mixture of Casodex<sup>®</sup> (bicalutamide) and/or its derivatives.

According to embodiments of the present invention, methods of preparing a  
 10 substantially pure enantiomer of an acylanilide such as Casodex<sup>®</sup> (bicalutamide) and/or its derivatives are provided. The methods include resolving an intermediate compound having the structure of Formula I:




The resolved intermediate compound of Formula I is then treated under conditions sufficient  
 15 to provide a substantially pure enantiomer of the acylanilide. Preferably, R<sup>1</sup> is lower alkyl having up to 4 carbons, and R<sup>2</sup> is lower alkyl having up to 6 carbons. More preferably, R<sup>1</sup> is methyl and R<sup>2</sup> is methylene. R<sup>3</sup> is preferably a direct link (*i.e.*, one or more bonds between X<sup>1</sup> and R<sup>4</sup>). R<sup>4</sup> is preferably phenyl which bears one, two or three substituents independently selected from hydrogen, halogen, nitro, carboxy, carbamoyl and cyano, and alkyl, alkoxy,  
 20 alkanoyl, alkylthio, alkylsulphinyl, alkylsulphonyl, perfluoroalkyl, perfluoroalkylthio, perfluoroalkylsulphinyl, perfluoroalkylsulphonyl, alkoxycarbonyl and N-alkylcarbamoyl each of up to 4 carbon atoms, and phenyl, phenylthio, phenylsulphinyl and phenylsulphonyl. More preferably, R<sup>4</sup> is phenyl which bears one, two or three substituents independently selected from hydrogen and halogen. Most preferably, R<sup>4</sup> is 4-fluorophenyl. Preferably, X<sup>1</sup>  
 25 is sulfur, sulphinyl, sulphonyl or imino. X<sup>1</sup> is more preferably sulfur, sulphinyl, or sulphonyl and is most preferably sulfur. The intermediate compounds according to the present invention may be prepared by various methods as will be understood by those skilled in the art. Exemplary methods of preparing particularly preferred intermediate compounds are described hereinbelow.

According to preferred embodiments of the present invention, a racemic mixture of the intermediate compound of Formula I is crystallizationally resolved. The intermediate compound of Formula I is contacted with a chiral base to obtain a diastereomeric mixture of chiral salts. The chiral salts of the diastereomeric mixture are then selectively crystallized to resolve the chiral salts into (R)- and (S)-salt enantiomers. The more preferred salt enantiomer is then treated to recover the more preferred hydroxyacid.

The chiral base is preferably (-)-cinchonidine. More preferably, the chiral base is (-)-cinchonidine (96%), which is commercially available from Aldrich of Milwaukee, Wisconsin. While the chiral base is preferably (-)-cinchonidine, other chiral bases may be employed including, but not limited to, brucine.

In particularly preferred embodiments, the substantially pure enantiomer of the more preferred (R)-hydroxyacid is resolved as follows:



The selective crystallization is preferably performed by contacting the diastereomeric mixture of the (R,-)-salt and the (S,-)-salt with a solvent system. The solvent system preferably includes methylene chloride and diethyl ether. The methylene chloride/diethyl ether solvent system preferably includes between about 1 and 40 percent by volume methylene chloride and between about 60 and 99 percent by volume diethyl ether, more preferably includes between about 5 and 30 percent by volume methylene chloride and between about 70 and 95 percent by volume diethyl ether, and most preferably includes between about 10 and 20 percent by volume methylene chloride and between about 80 and about 90 percent by volume diethyl ether. A particularly preferred solvent system consists essentially of between about


10 and 20 percent by volume methylene chloride and between about 80 and 90 percent by volume diethyl ether.

Progress of the resolution may be monitored by  $^{19}\text{F}$  NMR analysis, as illustrated in Figures 1 through 3. Figure 1 shows a  $^{19}\text{F}$  NMR spectrum of the pure salt of the (R)-hydroxyacid and (-)-cinchonidine. This (R)-hydroxyacid was made by asymmetric synthesis as described in the co-pending and co-assigned application entitled "Methods of Asymmetrically Synthesizing Enantiomers of Casodex, Its Derivatives and Intermediates Thereof" to Nnochiri N. Ekwuribe. This spectrum was used as a standard for comparison in determining the resolution of the racemic mixture. Figure 2 shows a  $^{19}\text{F}$  NMR spectrum of the salt of the racemic (R,S)-hydroxyacid and (-)-cinchonidine prior to resolution. The left portion of the signal is from the (S)-hydroxyacid and the right portion of the signal is from the (R)-hydroxyacid. Figure 3 shows a  $^{19}\text{F}$  NMR spectrum of the salt of the racemic (R,S)-hydroxyacid and (-)-cinchonidine after one round of crystallization. This spectrum illustrates that approximately 78% of the crystals formed were the cinchonidine salt of the (S)-hydroxyacid. Thus, the solution was becoming enriched with the salt of the more preferred (R)-hydroxyacid.


While those skilled in the art will understand how to recover the intermediate compound from the salt, the (R)-intermediate compound (e.g., (R)-hydroxyacid) is preferably recovered from the salt by acidification with aqueous 1M HCl followed by extraction with an organic solvent, such as methylene chloride or ethyl acetate.

According to another aspect of the present invention, the intermediate compound of Formula I may be separated by various means for physico-chemical separation known in the art, such as chromatographic resolution. Exemplary methods may be found in G. Subramanian, *A Practical Approach to Chiral Separations by Liquid Chromatography*, John Wiley & Sons, 1994; Thomas E. Beesley, Raymond P. W. Scoff, *Chiral Chromatography*, John Wiley & Son Ltd., 1999; and Satinder Ahuja, *Chiral Separations: Applications and Technology*, American Chemical Society, 1996. In a preferred method, the intermediates are separated using high pressure liquid chromatography (HPLC) by methods such as those described in Krstulovic, A.M., ed. *Chiral Separations by HPLC: Applications to Pharmacological Compounds*, Halsted Press, 1989. One skilled in the art will understand how to chromatographically resolve the intermediate compounds described herein.

In preferred embodiments, the intermediate compound has the structure of Formula III:




wherein X<sup>1</sup> is sulfur or sulphonyl. The intermediate compounds of Formula III may be prepared by various methods as will be understood by those skilled in the art. For example, the intermediate compounds of Formula III may be prepared according to methods described in Howard Tucker et al., *Resolution of the Nonsteroidal Antiandrogen 4'-Cyano-3-[(4-fluorophenyl)sulfonyl]-2-hydroxy-2-methyl-3'-(trifluoromethyl)-propioanilide and the Determination of the Absolute Configuration of the Active Enantiomer*, 31 J. MED. CHEM. 885-887 (1988), the disclosure of which is incorporated herein in its entirety. As another example, the intermediate compounds of Formula III may be prepared from the compound of Formula IV:




wherein X<sup>1</sup> is sulfur or sulphonyl. Hydrolysis of the compound of Formula IV will yield the compound of Formula III. When X<sup>1</sup> is sulfur, the compound may further be oxidized to provide a compound where X<sup>1</sup> is sulphonyl.

As still another example, the intermediate compounds of Formula III may be prepared using commercially available compounds as follows:




wherein  $\text{X}^2$  is a leaving group, preferably chloro, bromo, or iodo; NaH is sodium hydride; THF is tetrahydrofuran; and [O] is an oxidizing agent as will be understood by those skilled in the art.

- 5 As yet another example, the intermediate compound of Formula III may be prepared using commercially available materials as follows:



wherein mCPBA is meta-chloroperbenzoic acid, and hydrolysis is preferably performed using an aqueous acid or aqueous base solution. A racemic mixture of the intermediate compound of Formula III may be resolved according to methods of the present invention described above.

Once the more preferred substantially pure enantiomer of the intermediate compound of Formula I or III has been prepared by resolution as described above, the compound may be contacted with the aniline compound of Formula II:



to provide the more preferred substantially pure enantiomer of the acylanilide. R<sup>6</sup> is preferably cyano, fluoro, chloro, bromo, iodo, or hydrogen. More preferably, R<sup>6</sup> is cyano, fluoro, chloro, bromo, iodo, and, most preferably, R<sup>6</sup> is cyano. R<sup>7</sup> is preferably perfluoroalkyl, perfluoroalkylthio, perfluoroalkylsulphinyl or perfluoroalkylsulphonyl each of having up to 4 carbon atoms. More preferably, R<sup>7</sup> is perfluoroalkyl, and, most preferably, R<sup>7</sup> is perfluoromethyl. Most preferably, R<sup>8</sup> is hydrogen.

By resolving the racemic mixture of the intermediate compounds prior to synthesizing the intermediate with the more expensive aniline compound, methods according to the present invention may provide more cost effective ways of preparing substantially pure enantiomers of acylanilides such as Casodex<sup>®</sup> (bicalutamide) and/or derivatives thereof.

The present invention will now be described with reference to the following example. It should be appreciated that this example is for the purposes of illustrating aspects of the present invention, and does not limit the scope of the invention as defined by the claims.

15

### Example

#### **Formation of the salt**

The salt of the (R,S)-hydroxyacid and (-)-cinchonidine was formed by mixing a solution of the hydroxyacid (one equivalent) with a solution of (-)-cinchonidine (one equivalent). Cinchonidine was generally dissolved in chloroform, but the hydroxyacid may have been dissolved in solvents such as chloroform, methylene chloride, ethyl acetate, or ethanol. The mixture was then stirred overnight at room temperature. After removal of the solvent by rotary evaporation, the residual salt was then redissolved in the desired recrystallization solvent.

25

#### **Recrystallization**

A typical procedure involved placing 20 mg of the salt of the (R,S)-hydroxyacid and (-)-cinchonidine in a vial. Ethyl ether (2 mL) was then added. Methylene chloride was then added dropwise (with occasional shaking) just until the salt had been solubilized. The solution was then placed at 4°C for crystallization.

30

**Assaying resolution**

After crystallization had occurred, the supernatant was removed. The crystals were then dissolved in deuterated chloroform ( $\text{CDCl}_3$ ). The ratio of (R)- to (S)-enantiomer could then be assayed by integration of the fluorine signal(s) using  $^{19}\text{F}$  NMR.

5


In the drawings and specification, there have been disclosed typical preferred embodiments of the invention and, although specific terms are employed, they are used in a generic and descriptive sense only and not for purposes of limitation, the scope of the invention being set forth in the following claims.



**What Is Claimed Is:**

1. A method of preparing a substantially pure enantiomer of an acylanilide comprising:

(a) resolving an intermediate compound having the structure of Formula I:



5 wherein

$\text{R}^1$  is alkyl or haloalkyl having up to 4 carbons;

$\text{R}^2$  is alkyl having up to 6 carbon atoms;

$\text{R}^3$  is a direct link or alkyl having up to 6 carbon atoms;

10  $\text{R}^4$  is alkyl, alkenyl, hydroxyalkyl or cycloalkyl each of up to 6 carbons; or  $\text{R}^4$  is phenyl which bears one, two or three substituents independently selected from hydrogen, halogen, nitro, carboxy, carbamoyl and cyano, and alkyl, alkoxy, alkanoyl, alkylthio, alkylsulphinyl, alkylsulphonyl, perfluoroalkyl, perfluoroalkylthio, perfluoroalkylsulphinyl, perfluoroalkylsulphonyl, alkoxycarbonyl and N-alkylcarbamoyl each of up to 4 carbon atoms, and phenyl, phenylthio, phenylsulphinyl and phenylsulphonyl; or  $\text{R}^4$  is naphthyl; or  $\text{R}^4$  is a 5- or 6-membered saturated or unsaturated heterocyclic which contains one, two or three heteroatoms selected from oxygen, nitrogen and sulfur, which heterocyclic may be a single ring or may be fused to a benzo-ring, and which heterocyclic is unsubstituted or bears one or two halogen, cyano or amino, or alkyl, alkoxy, alkylthio, alkylsulphinyl or alkylsulphonyl each of up to 4 carbon atoms, or oxy or hydroxy substituents, or which if sufficiently saturated may bear one or two oxo substituents; and

15  $\text{X}^1$  is oxygen, sulfur, sulphinyl (-SO-), sulphonyl (-SO<sub>2</sub>-), imino (-NH-) or alkylimino (-NR<sup>5</sup>-) where  $\text{R}^5$  is alkyl having up to 6 carbon atoms; and

20 (b) treating the resolved intermediate compound of Formula I under conditions sufficient to provide a substantially pure enantiomer of an acylanilide.

2. The method according to Claim 1, wherein the resolution step comprises crystallizationally resolving the intermediate compound of Formula I.

3. The method according to Claim 2, wherein the crystallizationally resolving step comprises:

contacting the intermediate compound of Formula I with a chiral base to provide a diastereomeric mixture of a chiral salt;

5 crystallizationally resolving the diastereomeric mixture of the chiral salt to provide a substantially pure enantiomer of the chiral salt; and

recovering a substantially pure enantiomer of the intermediate compound of Formula I.

4. The method according to Claim 3, wherein the contacting step comprises contacting the intermediate compound of Formula I with (-)-cinchonidine to provide a diastereomeric mixture of intermediate compound-cinchonidine chiral salts.

5. The method according to Claim 3, wherein the step of crystallizationally resolving the diastereomeric mixture of the chiral salt comprises contacting the diastereomeric mixture of the chiral salt with a solvent system comprising methylene chloride and diethyl ether.

6. The method according to Claim 5, wherein the solvent system comprises between about 1 and 40 percent by volume methylene chloride and between about 60 and 99 percent by volume diethyl ether.

7. The method according to Claim 6, wherein the solvent system consists essentially of between about 10 and 20 percent by volume methylene chloride and between about 80 and 90 percent by volume diethyl ether.

8. The method according to Claim 1, wherein  $R^1$  is alkyl having up to 4 carbons;  $R^2$  is alkyl having up to 6 carbons;  $R^3$  is a direct link;  $R^4$  is phenyl having one, two or three substituents independently selected from hydrogen, halogen, nitro, carboxy, carbamoyl and cyano, and alkyl, alkoxy, alkanoyl, alkylthio, alkylsulphinyl, alkylsulphonyl, perfluoroalkyl, perfluoroalkylthio, perfluoroalkylsulphinyl, perfluoroalkylsulphonyl, alkoxycarbonyl and N-


5

alkylcarbamoyl each of up to 4 carbon atoms, and phenyl, phenylthio, phenylsulphinyl and phenylsulphonyl; and  $X^1$  is sulfur, sulphinyl, sulphonyl or imino.

9. The method according to Claim 1, wherein  $R^1$  is methyl,  $R^2$  is methylene,  $R^3$  is a direct link,  $R^4$  is 4-fluorophenyl, and  $X^1$  is sulfur, sulphinyl, or sulphonyl.

10. The method according to Claim 1, wherein the resolution step comprises high performance liquid chromatographically resolving the intermediate compound of Formula I.

11. The method according to Claim 1, wherein step (b) comprises contacting the resolved intermediate with an aniline compound having the structure of Formula II:



Formula II

wherein

5  $R^6$  is cyano, carbamoyl, nitro, fluoro, chloro, bromo, iodo, or hydrogen, or alkyl, alkoxy, alkanoyl, alkylthio, alkylsulphinyl, alkylsulphonyl, perfluoroalkyl, perfluoroalkylthio, perfluoroalkylsulphinyl or perfluoroalkylsulphonyl each having up to 4 carbon atoms, or phenylthio, phenylsulphinyl or phenylsulphonyl;

10  $R^7$  is cyano, carbamoyl, nitro, fluoro, chloro, bromo or iodo, or alkanoyl, alkylthio, alkylsulphinyl, alkylsulphonyl, perfluoroalkyl, perfluoroalkylthio, perfluoroalkylsulphinyl or perfluoroalkylsulphonyl each of having up to 4 carbon atoms; or phenylthio, phenylsulphinyl or phenylsulphonyl; and


$R^8$  is hydrogen or halogen;

under conditions sufficient to provide a substantially pure enantiomer of an acylanilide.

12. The method according to Claim 11, wherein  $R^6$  is cyano,  $R^7$  is perfluoromethyl, and  $R^8$  is hydrogen.

13. A method of preparing a substantially pure enantiomer of Casodex<sup>®</sup> (bicalutamide) or a derivative thereof comprising:

(a) resolving an intermediate compound having the structure of Formula I:



5 wherein

$\text{R}^1$  is methyl;

$\text{R}^2$  is methylene;

$\text{R}^3$  is a direct link;

$\text{R}^4$  is 4-fluorophenyl; and

10  $\text{X}^1$  is sulfur, sulphinyl ( $-\text{SO}-$ ), or sulphonyl ( $-\text{SO}_2-$ ); and

(b) treating the resolved intermediate compound of Formula I under conditions sufficient to provide a substantially pure enantiomer of Casodex<sup>®</sup> (bicalutamide) or a derivative thereof.

14. The method according to Claim 1, wherein the resolution step comprises crystallizationally resolving the intermediate compound of Formula I.

15. The method according to Claim 14, wherein the crystallizationally resolving step comprises:

contacting the intermediate compound of Formula I with a chiral base to provide a diastereomeric mixture of a chiral salt;

5 crystallizationally resolving the diastereomeric mixture of the chiral salt to provide a substantially pure enantiomer of the chiral salt; and

recovering a substantially pure enantiomer of the intermediate compound of Formula I.

16. The method according to Claim 15, wherein the contacting step comprises contacting the intermediate compound of Formula I with (-)-cinchonidine to provide a diastereomeric mixture of intermediate compound-cinchonidine chiral salts.


17. The method according to Claim 15, wherein the step of crystallizationally resolving the diastereomeric mixture of the chiral salt comprises contacting the

diastereomeric mixture of the chiral salt with a solvent system comprising methylene chloride and diethyl ether.

18. The method according to Claim 17, wherein the solvent system comprises between about 1 and 40 percent by volume methylene chloride and between about 60 and 99 percent by volume diethyl ether.

19. The method according to Claim 17, wherein the solvent system consists essentially of between about 10 and 20 percent by volume methylene chloride and between about 80 and 90 percent by volume diethyl ether.

20. The method according to Claim 13, wherein step (b) comprises contacting the resolved intermediate with an aniline compound having the structure of Formula II:



Formula II


wherein


5      R<sup>6</sup> is cyano;


        R<sup>7</sup> is perfluoroalkyl; and

        R<sup>8</sup> is hydrogen;

under conditions sufficient to provide a substantially pure enantiomer of Casodex<sup>®</sup> (bicalutamide).

**Figure 1**

**Figure 2**



**Figure 3**



# INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/US 00/41609

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 C07C315/06 C07C317/28

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data, BEILSTEIN Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 95 19770 A (SEPRACOR INC) 27 July 1995 (1995-07-27) cited in the application page 9, line 8-26 ---	1-20
Y	WO 94 08986 A (PFIZER ;MURTIASHAW CHARLES W III (US); VANDERPLAS BRIAN C (US)) 28 April 1994 (1994-04-28) the whole document --- -/--	1-20

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

4 April 2001

Date of mailing of the international search report

12/04/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Goetz, G

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/41609

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>TUCKER H ET AL: "RESOLUTION OF THE NONSTEROIDAL ANTIANDROGEN 4'-CYANO-3-U(4-FLUOROPHENYL)SULFONYL-2-HYDROXY-2METHYL-3'-(TRIFLUOROMETHYL)- PROPIONANILIDE AND THE DETERMINATION OF THE ABSOLUTE CONFIGURATION OF THE ACTIVE ENANTIOMER" JOURNAL OF MEDICINAL CHEMISTRY,US,AMERICAN CHEMICAL SOCIETY. WASHINGTON, vol. 31, no. 4, 1 April 1988 (1988-04-01), pages 885-887, XP000605163  ISSN: 0022-2623  cited in the application  the whole document  -----</p>	1-20

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/41609

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9519770 A	27-07-1995	AU 1687395 A	08-08-1995
		AU 3507499 A	19-08-1999
		CA 2181358 A	27-07-1995
		EP 0748220 A	18-12-1996
		JP 9508125 T	19-08-1997
		US 5985868 A	16-11-1999
WO 9408986 A	28-04-1994	AU 4650493 A	09-05-1994
		CA 2146005 A	28-04-1994
		CN 1090577 A	10-08-1994
		CZ 9501008 A	18-10-1995
		EP 0665839 A	09-08-1995
		FI 934624 A	22-04-1994
		HU 65128 A	28-04-1994
		JP 7507811 T	31-08-1995
		MX 9306526 A	29-04-1994
		NO 951507 A	20-04-1995
		PL 308473 A	07-08-1995
		ZA 9307737 A	19-04-1995